



UNIVERSITI PUTRA MALAYSIA

**ISOLATION AND BIOLOGICAL ACTIVITY OF NATURALLY
OCCURRING COMPOUNDS FROM MELICOPE LUNU ANKENDA
(GAERTN) T-HARTLEY, MELICOPE BONWICKII(F-MUELL) T-
HARYLEY AND TETRADIUM SAMBUCINUM (BL) HARTLEY**

ISMIARNI KOMALA.

FS 2005 30

**ISOLATION AND BIOLOGICAL ACTIVITY OF NATURALLY
OCCURRING COMPOUNDS FROM *MELICOPE LUNU ANKENDA*
(GAERTN) T-HARTLEY, *MELICOPE BONWICKII* (F-MUELL)
T-HARTLEY AND *TETRADIUM SAMBUCINUM* (BL) HARTLEY**

By

ISMIARNI KOMALA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for Degree of Master of Science**

June 2005



Abstract of thesis presented to Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**ISOLATION AND BIOLOGICAL ACTIVITY OF NATURALLY
OCCURRING COMPOUNDS FROM *MELICOPE LUNU ANKENDA*
(GAERTN) T-HARTLEY, *MELICOPE BONWICKII* (F-MUELL)
T-HARTLEY AND *TETRADIUM SAMBUCINUM* (BL) HARTLEY**

By

ISMIARNI KOMALA

June 2005

Chairman : Professor Mawardi Rahmani, PhD

Faculty : Science

Detailed investigation on three Rutaceous plants have resulted in the isolation and structural elucidation of a number of compounds. The structures of these compounds were elucidated by using spectroscopic methods such as UV (ultraviolet), IR (infra red), MS (mass spectra), NMR (Nuclear Magnetic Resonance) and also by comparison with previous reports. The crude extracts and isolated compounds were evaluated for their antioxidant, cytotoxicity and antimicrobial activities using DPPH (1,1-diphenil-2-picrylhydrazyl), MTT ((3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) and disc diffusion methods, respectively.

The phytochemical study on leaves of *Melicope lunu-ankenda* (Gaertn) T. Hartley afforded *p*-*O*-geranylcoumaric acid (153) sesamin (109), melisemine (154) and a new compound given tentative structure of 7,7''-digeranyloxy-2'',3''-epoxycinnamic anhydride (155). The melisemine (154) showed weak activity as a radical scavenger with an IC₅₀ value of 75 µg/mL. Meanwhile the 7,7''-digeranyloxy-2'',3''-

epoxycinnamic anhydride (155) was found to be active against cervical cancer (Hela) cell line with an IC_{50} value of 34 $\mu\text{g/mL}$.

The investigations on leaves of *Melicope bonwickii* (F. Muell) T. Hartley have resulted in the isolation of three known furoquinoline alkaloid, 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156), evellerine (157) and kokusaginine (11) together with a new 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158) and a known amide compound aurantiamide acetate (159). Two compounds 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156) and 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158) were found to be toxic to cervical cancer (Hela) cell line with IC_{50} values of 6.0 and 11.4 $\mu\text{g/mL}$, respectively.

Detailed extraction and separation of bark and leaves of *Tetradium sambucinum* (Bl) Hartley have led to the isolation of decarine (160), rutaecarpine (122) 7-hydroxycoumarin (141) and aurantiamide acetate (159). Decarine (160) was active against the cervical cancer cell lines (Hela) with an IC_{50} value of 14.6 $\mu\text{g/mL}$, while rutaecarpine (122) showed weak a radical scavenger activity with an IC_{50} value of 75 $\mu\text{g/mL}$.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan ijazah Master Sains

**PEMENCILAN DAN AKTIVITI BIOLOGI SEBATIAN SEMULAJADI
DARIPADA *MELICOPE LUNU ANKENDA* (GAERTN) T-HARLEY,
MELICOPE BONWICKII (F. MUELL) T-HARTLEY DAN *TETRADIUM
SAMBUCINUM* (BL) HARTLEY**

Oleh

ISMIARNI KOMALA

Jun 2005

Pengerusi : Professor Mawardi Rahmani, PhD

Fakulti : Sains

Kajian terhadap tiga tumbuhan dalam famili Rutaceae telah menghasilkan pemencilan beberapa sebatian. Struktur dari sebatian-sebatian ini dikenal pasti dengan menggunakan kaedah spektroskopi seperti IR, UV, NMR, MS dan juga perbandingan dengan kajian-kajian lepas. Ekstrak mentah dan sebatian-sebatian yang telah dipencilkan dari tumbuhan ini diuji untuk aktiviti antioksidan, sitotoksik dan antimikrob dengan menggunakan kaedah DPPH (1,1-diphenil-2-picrylhydrazyl), MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) dan peresapan cakera.

Kajian fitokimia ke atas daun *Melicope lunu-ankenda* (Gaertn) T. Hartley telah menghasilkan dua sebatian yang telah dikenali sebagai *p*-*O*-geranylecoumaric acid (153) sesamin (109) melisemine (154) dan struktur sementara daripada 7,7'-digeranyloxy-2'',3''-epoxycinnamic anhydride (155). Melisemine (154) telah menunjukkan aktiviti yang lemah sebagai antioksidan dengan nilai IC₅₀ 75 µg/mL.

Manakala sebatian 7,7''-digeranyloxy-2'',3''-epoxycinnamic anhydride (155) telah menunjukkan kesan aktiviti terhadap kanser rahim (Hela) dengan nilai IC_{50} 34 $\mu\text{g/mL}$.

Pengkajian terhadap daun *Melicope bonwickii* (F. Muell) T. Hartley telah menghasilkan 3 alkaloid furoquinoline yang telah dikenali: 7-(2',3'-epoxypronyloxy)-4-methoxyfuroquinoline (156) evellerine (157) dan kokusaginine (11) bersama dengan sebatian baru 7-(2-hydroxy-3'-chloropronyloxy)-4-methoxyfuroquinoline (160) dan sebatian amide yang telah dikenali, aurantiamide acetate (159). Dua sebatian yang telah dipencilkan daripada daun *Melicope bonwickii* (F. Muell) T. Hartley yaitu 7-(2',3'-epoxypronyloxy)-4-methoxyfuroquinoline (158) dan 7-(2-hydroxy-3'-chloropronyloxy)-4-methoxyfuroquinoline (160) telah diujikan aktiviti dengan sel kanser rahim (Hela), hasilnya menunjukkan kedua-dua sebatian ini mempunyai kesan sitotoksik yang masing-masing mempunyai nilai IC_{50} 6.0 dan 11.4 $\mu\text{g/mL}$.

Pengestrakan dan pemisahan daripada kulit batang dan daun *Tetradium sambucinum* (Bl) Hartley telah membawa kepada pemencilan sebatian-sebatian decarine (160), rutaecarpine (122) 7-hydroxycoumarin (141) dan aurantiamide acetate (159). Decarine menunjukkan aktiviti yang sederhana terhadap sel kanser rahim (Hela) yang mempunyai nilai IC_{50} 34 $\mu\text{g/mL}$, manakala rutaecarpine (122) menunjukkan aktiviti yang lemah sebagai antioksidan dengan nilai IC_{50} 75 $\mu\text{g/mL}$.

ACKNOWLEDGEMENTS

All praises do to Allah, Lord of the universe. Only by his grace and mercy this thesis can be completed.

I wish to express my sincere thanks to my supervisor Prof. Dr. Mawardi Rahmani for his invaluable guidance, support and continuous encouragement throughout the course of this project.

My gratitude also goes to the members of my supervisory committee, Assoc. Prof. Dr. Gwendoline Ee Cheng Lian and Assoc. Prof. Dr. Asmah Rahmat for their assistance in my research.

Thanks are also due to Prof. Dr. Dachriyanus, Apt, Andalas University, Indonesia, for suggestions in my research and for helping the plant collection from Indonesia, Drs. Rusdji Tamin, Andalas University Herbarium for identifying the plant material.

Financial support the Malaysian Government IRPA program is gratefully acknowledged

Special thanks are extended to other members of the academic and technical staff, students and friends who have helped me in every way possible.

Finally, my deepest thanks to my parents, sister and brother for their prayers, continuous moral support and unending encouragement.

I certify that an Examination Committee met on 20th June 2005 to conduct the final examination of Ismiarni Komala on her Master of Science thesis entitled “Isolation and Biological Activity of Naturally Occuring Compounds from *Melicope lunu ankenda* (Gaertn) T-Hartley, *Melicope bonwickii* (F-Muell) T-Hartley and *Tetradium sambucinum* (BL) Hartley” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Nordin Lajis, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Mohd. Aspollah Md. Sukari, PhD


Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Khozirah Shaari, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Dato' Laily Din PhD

Professor
Centre for Graduate Studies
Universiti Kebangsaan Malaysia
(External Examiner)



GULAM RUSUL RAHMAT ALI, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 JUL 2005



This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of Supervisory Committee are as follow:

Mawardi Rahmani, PhD


Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Gwendoline Ee Cheng Lian, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Asmah Rahmat, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Member)



AINI IDERIS, PhD

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 11 AUG 2005

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledgment. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



ISMIARNI KOMALA

Date **29/7/2005**

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vi
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xx
CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	5
The genus <i>Melicope</i>	5
Distribution and General Information	5
Medicinal Uses	5
<i>Melicope lumu-ankenda</i> (Gaertn) T. Harley	6
<i>Melicope bonwickii</i> (F.Muell) T. Hartley	7
<i>Tetradium sambucinum</i> (Bl) Hartley	7
Chemical Constituent and Biological Activity	8
Bioassay	31
Antioxidant Activity Assay	31
Cytotoxic Activity Assay	33
Antimicrobial Activity assay	34
III MATERIALS AND METHODS	36
General Experimental Methods	36
Materials and Extraction	36
Instrument	36
Chromatography	38
Extraction and Isolation of Compounds from Leaves of <i>Melicope lumu-ankenda</i> (Gaertn) T.Harley	39
Extraction and Separation	39
Extraction and Isolation of Compound from Leaves of <i>Melicope bonwickii</i> (F.Muell) T.Hartley	44
Extraction and Separation	44
Extraction and Isolation of Compounds from <i>Tetradium sambucinum</i> (Bl) Hartley	50
Extraction and Separation of Barks	50
Extraction and Separation of Leaves	51
Bioassay	55
Antioxidant Activity Assay	55
Cytotoxic Activity Assay	56
Antimimicrobial Activity Assay	57



IV	RESULTS AND DISCUSSIONS	59
	Characterization of the Isolated Compounds from Leaves of <i>Melicope lunu-ankenda</i> (Gaertn) T.Harley	59
	<i>p</i> - <i>O</i> -Geranylcoumaric acid(153)	60
	Sesamin (109)	73
	Melisemin (154)	85
	7,7''-Digeranyloxy-2'',3''-epoxycinnamic anhydride (155)	97
	Characterization of Isolated Compounds from Leaves of <i>Melicope bonwickii</i> (F.Muell) T.Hartley	112
	7-(2',3'-Epoxyprenyloxy)-4-methoxyfuroquinoline (156)	112
	Evellerine (157)	126
	7-(2'-Hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158)	139
	Aurantiamide acetate (159)	155
	Kokusaginine (11)	170
	Characterization of Isolated Compounds from Barks of <i>Tetradium sambucinum</i> (Bl) Hartley	177
	Decarine (160)	177
	Characterization of Isolated Compounds from Leaves of <i>Tetradium sambucinum</i> (Bl) Hartley	190
	Rutaecarpine (122)	190
	7-Hydroxycoumarin (141)	208
	Aurantiamide acetate (159)	216
	Bioassay	217
	Antioxidant Activity Assay	217
	Cytotoxic Activity Assay	221
	Antimicrobial Activity Assay	224
V	CONCLUSIONS	227
	REFERENCES	231
	BIODATA OF THE AUTHOR	

LIST OF TABLES

Table	Page
1. NMR spectral data of <i>p</i> - <i>O</i> -geranylcoumaric acid (153)	65
2. NMR spectral data of sesamin (109)	78
3. NMR spectral data of melisemine (154)	90
4. NMR spectral data of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	102
5. NMR spectral data 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	118
6. NMR spectral data of evellerine (157)	131
7. NMR spectral data of 7-(2-hydroxy-3-chloroprenyloxy)-4-methoxyfuroquinoline (158)	145
8. NMR spectral data of aurantiamide acetate (159)	161
9. NMR spectral data of kokusaginine (11)	173
10. NMR spectral data of decarine (160)	182
11. NMR spectral of rutaecarpine (122)	197
12. NMR spectral data of 7-hydroxycoumarin (141)	211
13. Radical Scavenging activity of crude extracts and isolated compounds from <i>Melicope lunu-ankenda</i> (Gaertn) T.Hartley <i>Evodia sambucina</i> Hook. f. and <i>Melicope bonwicckii</i> (F.Muell) T.Hartley.	218
14. Cytotoxic activity of crude extracts and isolated compounds from <i>Melicope lunu-ankenda</i> (Gaertn) T.Hartley <i>Evodia sambucina</i> Hook. f. and <i>Melicope bonwickii</i> (F.Muell) T.Hartley.	223
15. Antimicrobial activity of crude extracts and isolated compounds from <i>Melicope lunu-ankenda</i> (Gaertn) T.Hartley <i>Evodia sambucina</i> Hook. f. and <i>Melicope bonwickii</i> (F.Muell), T.Hartley	226



LIST OF FIGURES

Figure	Page
1. Metabolism of MTT to a farmazan by viable cells	34
2. UV spectrum of <i>p</i> - <i>O</i> -Geranylcoumaric acid (153)	60
3. IR spectrum of <i>p</i> - <i>O</i> -Geranylcoumaric acid (153)	61
4. Selectected HMBC correlations of <i>p</i> - <i>O</i> -Geranylcoumaric acid (153)	63
5. Mass fragmentation patterns of <i>p</i> - <i>O</i> -Geranylcoumaric acid (153)	64
6. Mass spectrum of <i>p</i> - <i>O</i> -geranylcoumaric acid (153)	66
7. ¹ H-NMR spectrum of <i>p</i> - <i>O</i> -geranylcoumaric acid (153)	67
8. ¹³ C-NMR spectrum of <i>p</i> - <i>O</i> -geranylcoumaric acid (153)	68
9. DEPT spectrum of <i>p</i> - <i>O</i> -geranylcoumaric acid (153)	69
10. COSY spectrum of <i>p</i> - <i>O</i> -geranylcoumaric acid (153)	70
11. HSQC spectrum of <i>p</i> - <i>O</i> -geranylcoumaric acid (153)	71
12. HMBC spectrum of <i>p</i> - <i>O</i> -geranylcoumaric acid (153)	72
13. UV spectrum of sesamin (109)	73
14. IR spectrum of sesamin (109)	74
15. Selected HMBC correlations of sesamin	76
16. Mass fragmentation patterns of sesamin (109)	77
17. Mass spectrum of sesamin (109)	79
18. ¹ H-NMR spectrum of sesamin (109)	80
19. ¹³ C-NMR spectrum of sesamin (109)	81
20. COSY spectrum of sesamin (109)	82
21. HSQC spectrum of sesamin (109)	83
22. HMBC spectrum of sesamin (109)	84

23. UV spectrum of melisemine (154)	86
24. IR spectrum of melisemine (154)	86
25. Selected HMBC correlations of melisemine (154)	88
26. Mass fragmentation patterns of melisemine (154)	89
27. Mass spectrum of melisemine (154)	91
28. ¹ H-NMR spectrum of melisemine (154)	92
29. ¹³ C-NMR spectrum of melisemine (154)	93
30. COSY spectrum of melisemine (154)	94
31. HSQC spectrum of melisemine (154)	95
32. HMBC spectrum of melisemine (154)	96
33. UV spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	98
34. IR spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	98
35. Selected HMBC correlations of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	101
36. EIMS spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	103
37. FABMS spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	104
38. HRFABMS spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	105
39. ¹ H-NMR spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	106
40. ¹³ C-NMR spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	107
41. DEPT spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	108
42. COSY spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	109
43. HSQC spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	110
44. HMBC spectrum of 7,7''-digeranyloxy-2',3'-epoxycinnamic anhydride (155)	111

45. UV spectrum of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	113
46. IR spectrum of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	113
47. HMBC correlations of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	116
48. Mass fragmentation patterns of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	117
49. Mass spectrum of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	119
50a. ¹ H-NMR spectrum of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	120
50b. Expanded ¹ H-NMR spectrum of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	121
50c. Expanded ¹ H-NMR spectrum of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	122
51. ¹³ C-NMR spectrum of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	123
52. COSY spectrum of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	124
53. HMBC spectrum of 7-(2',3'-epoxyprenyloxy)-4-methoxyfuroquinoline (156)	125
54. UV spectrum of evellerine (157)	127
55. IR spectrum evellerine (157)	127
56. Selected HMBC correlations spectrum of evellerine (157)	129
57. Mass fragmentation patterns of evellerine (157)	130
58. Mass spectrum of evellerine (157)	132
59a. ¹ H-NMR spectrum evellerine (157)	133
59b. Expanded ¹ H-NMR spectrum of evellerine (157)	134
59c. Expanded ¹ H-NMR spectrum of evellerine (157)	135
60. ¹³ C-NMR spectrum of evellerine (157)	136
61. COSY spectrum of evellerine (157)	137
62. HMBC spectrum of evellerine (157)	138

63. UV spectrum 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (157)	140
64. IR spectrum 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (157)	140
65. Selected HMBC correlations of 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158)	143
66. Mass fragmentation patterns of 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158)	144
67. Mass spectrum 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (157)	146
68a. ¹ H-NMR spectrum 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (157)	147
68b. Expanded ¹ H-NMR spectrum of 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158)	148
68c. Expanded ¹ H-NMR spectrum of 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline	149
69. ¹³ C-NMR spectrum of 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158)	150
70. COSY spectrum of 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158)	151
71. DEPT spectrum of 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158)	152
72. HSQC spectrum of 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158)	153
73. HMBC spectrum of 7-(2'-hydroxy-3'-chloroprenyloxy)-4-methoxyfuroquinoline (158)	154
74. UV spectrum of aurantiamide acetate (159)	155
75. IR spectrum of aurantiamide acetate (159)	156
76. Selected HMBC correlations of aurantiamide acetate (159)	158
77. Mass Fragmentation patterns of aurantiamide acetate (159)	160
78. Mass spectrum of aurantiamide acetate (159)	162
79a. ¹ H-NMR spectrum of aurantiamide acetate (159)	163

79b. Expanded ^1H -NMR spectrum of aurantiamide acetate (159)	164
80a. ^{13}C -NMR spectrum of aurantiamide acetate (159)	165
80b. Expanded ^{13}C -NMR spectrum of aurantiamide acetate (159)	166
81. COSY spectrum of aurantiamide acetate (159)	167
82. HSQC spectrum of aurantiamide acetate (159)	168
83. HMBC spectrum of aurantiamide acetate (159)	169
84. UV spectrum of kokusaginine (11)	171
85. IR spectrum of kokusaginine (11)	171
86. Mass fragmentation patterns of kokusaginine (11)	172
87. Mass spectrum of kokusaginine (11)	174
88a. ^1H -NMR spectrum of kokusaginine (11)	175
88b. Expanded ^1H -NMR spectrum of kokusaginine (11)	176
89. UV spectrum of decarine (160)	178
90. IR spectrum of decarine (160)	178
91. Selected HMBC correlations of decarine (160)	180
92. Mass fragmentation patterns of decarine (160)	181
93. Mass spectrum of decarine (160)	183
94. ^1H -NMR spectrum of decarine (160)	184
95. ^{13}C -NMR spectrum of decarine (160)	185
96. COSY spectrum of decarine (160)	186
97. DEPT spectrum of decarine (160)	187
98. HSQC spectrum of decarine (160)	188
99. HMBC spectrum of decarine (160)	189
100. UV spectrum of rutaecarpine (122)	191
101. IR spectrum of rutaecarpine (122)	191

102. Selected HMBC correlations of rutaecarpine (122)	194
103a. Mass fragmentation patterns of rutaecarpine (122)	195
103b. Mass fragmentation patterns of rutaecarpine (122)	196
104. Mass spectrum of rutaecarpine (122)	198
105. ¹ H-NMR spectrum of rutaecarpine (122)	199
106a. ¹³ C-NMR spectrum of rutaecarpine (122)	200
106b. Expanded ¹³ C-NMR spectrum of of rutaecarpine (122)	201
107. COSY spectrum of rutaecarpine (122)	202
108. DEPT spectrum of rutaecarpine (122)	203
109. HSQC spectrum of rutaecarpine (122)	204
110a. HMBC spectrum of rutaecarpine (122)	205
110b. Expanded HMBC spectrum of rutaecarpine (122)	206
110c. Expanded HMBC spectrum of rutaecarpine (122)	207
111. UV spectrum of 7-hydroxycoumarin (141)	209
112. IR spectrum of 7-hydroxycoumarin (141)	209
113. Mass fragmentation patterns of 7-hydroxycoumarin (141)	210
114. Mass spectrum of 7-hydroxycoumarin (141)	212
115. ¹ H-NMR spectrum of 7-hydroxycoumarin (141)	213
116. ¹³ C-NMR spectrum of 7-hydroxycoumarin (141)	214
117. COSY spectrum of 7-hydroxycoumarin (141)	215
118. Radical scavenging activity of <i>Melicope lumu-ankenda</i> (Gaertn) T.Hartley leaves	219
119. Radical scavenging activity of barks of <i>Tetradium sambucinum</i> (Bl) Hartley	219
120. Radical scavenging activity of leaves of <i>Tetradium sambucinum</i> (Bl) Hartley	220

LIST OF ABBREVIATIONS

α	alpha
β	betha
δ	delta (chemical shift in ppm)
γ	gamma
λ_{max}	maximum wavelength in nm
μg	microgram
μl	microliter
Ar	aromatic
ATP	adenosine tri phosphate
br	broad
^{13}C	carbon-13
$^{\circ}\text{C}$	degree celcius
CFU	colony forming unit
CHCl_3	chloroform
CDCl_3	deuterated choroform
cm^{-1}	per centimeter
COSY	Correlated Spectroscopy
<i>d</i>	doublet
<i>dd</i>	doublet of doublets
<i>ddd</i>	doublet of doublets of doublets
DPPH	1,1-diphenil-2-picrylhydrazyl
DEPT	Distortionless Enhacement by Polarization Transfer
DMSO	Dimethylsulfoxides

EIMS	Electron Impact Mass Spectroscopy
ELISA	Enzym-Linked Imunosorbent Assay
FCS	Fetal Calf Serum
g	gram
GCMS	Gas Chromatography-Mass Spectroscopy
h	hour
^1H	proton
HMBC	Heteronuclear Multiple Bond Connectivity by 2D Multiple Quantum
HSQC	Heteronuclear Single Quantum Coherence
Hz	hertz
IC ₅₀	Inhibition Concentration at 50 percent
IR	Infra red
J	coupling constant value
KBr	Kalium bromida
Kg	kilogram
M ⁺	molecular ion
m	multiplet
mg	milligram
mL	milliliter
mm	millimeter
MeOH	methanol
MHz	MegaHertz
m.p.	melting point
MTT	Microculture tetrazolium
	3-[4,5-dimethylthiazol-2yl)-2,5-diphenyltetrazolium bromide

m/z	mass per charge
Na	Natrium
NA	Natrium agar
nm	nanometer
NMR	Nuclear Magnetic Resonance
<i>o</i>	ortho
OD	Optical density
<i>p</i>	para
PBS	Phosphate Buffered Saline
Pet.ether	Petroleum ether
ppm	part per million
RPMI	Roswell Park Memorial Institute
<i>s</i>	singlet
<i>t</i>	triplet
TLC	Thin Layer Chromatography
TMS	Tetramethilsilane
UV	ultra violet
WHO	World Health Organization.



CHAPTER I

INTRODUCTION

The use of natural products with therapeutic properties is as ancient as human civilization and for a long time, mineral, plant and animal products were the main sources of drug. Even today, plants are the almost exclusive source of drug for the majority of the world's populations.

About 25 % of drugs prescribed worldwide come from plants and 121 such active compounds are currently being used. Of the 252 drugs considered as basic and essential by the World Health Organization (WHO), 11% are exclusively of plant origin and a significant number are synthetic drugs obtained from natural precursors (Rates, 2001). This is particularly true as shown in cancer and infectious diseases, where over 60% and 75% of the drugs were known to be of natural origin, respectively (Newman *et al.*, 2003).

After centuries of empirical use of herbal preparation, the isolation of active principles such as morphine, strychnine, quinine etc. in the early 19th century marked a new era in the use of medicinal plants, and the beginning of modern medicinal plant research. Plant metabolites were mainly investigated from a phytochemical and chemotaxonomic viewpoint during this period. Over the last decade, however, interest in drugs of plant origin has been growing steadily. Consumption of medicinal plants has almost doubled in Western Europe during that period (Hamburger and Hostettman, 1991).



It is currently estimated that there are at least 250.000-500.000 different plants species, up to 30 millions species of insects, 1.2 millions species of fungi and similar number of algae and prokaryotes in existence throughout the world (Pimm *et al.*, 1995). All of the species coexist in ecosystem and interact with each other in several ways in which chemistry plays a major role, for example in defense, symbiosis and pollination. In basic term, these organisms all share a similar biochemical process necessary for living cell, but in addition to that they also produce a wide variety of the so called secondary metabolites that are involved in interactions between organisms. Considering the number of organisms, and the almost infinite number of interactions possible, it is not surprising that an enormously wide variety of secondary metabolites have evolved within organism (Vepoorte, 1998).

The potential of higher plants as sources for new drugs is still largely unexplored. Among the estimated 250.000-500.000 plants species, only a small percentage has been investigated phytochemically and the fraction submitted to biological or pharmacological screening is even smaller (Rates, 2001). Plants contain hundreds or thousands of metabolites. It is clear that plants provide an enormous potential for the discovery of new bioactive compounds.

Melicope species is one of the genus in Rutaceae family. *Melicope* have similar characteristic with *Tetradium* Lour and *Euodia* and recently Hartley reestablished the genus *Tetradium* Lour that was formerly placed under *Euodia* J.R. & G. Forst by Engler (Engler, 1931). Hartley further indicated that many other species formerly placed in *Euodia* were to be transferred to *Melicope* J.R & G. Forst. (Hartley, 1981)